

*Journal of Chromatography*, 379 (1986) 347–366  
*Biomedical Applications*  
 Elsevier Science Publishers B.V., Amsterdam — Printed in The Netherlands

CHROMBIO. 2992

REVIEW

GAS CHROMATOGRAPHIC PROFILING OF KETONE BODIES AND  
 ORGANIC ACIDS IN DIABETES\*

H.M. LIEBICH

*Medizinische Universitätsklinik, D-7400 Tübingen (F.R.G.)*

(First received November 26th, 1985; revised manuscript received December 29th, 1985)

CONTENTS

1. Introduction. . . . .	348
2. Profiling procedures. . . . .	349
2.1. Methods of choice. . . . .	349
2.2. Ketones. . . . .	349
2.2.1. Isolation of ketones. . . . .	349
2.2.2. Gas chromatographic profiling of ketones. . . . .	349
2.3. Organic acids. . . . .	350
2.3.1. Isolation of organic acids. . . . .	350
2.3.2. Derivatization of carboxyl groups. . . . .	350
2.3.3. Derivatization of carbonyl groups. . . . .	351
2.3.4. Pre-fractionation of organic acids. . . . .	351
2.3.5. Gas chromatographic profiling of organic acids. . . . .	352
3. Application to diabetes mellitus. . . . .	353
3.1. Diabetic patients under good therapeutic control. . . . .	353
3.2. Diabetic patients with ketoacidosis. . . . .	354
3.2.1. Ketones. . . . .	354
3.2.2. Fatty acids. . . . .	357
3.2.3. Aliphatic dicarboxylic acids. . . . .	357
3.2.4. Aliphatic oxocarboxylic acids. . . . .	359
3.2.5. Aliphatic hydroxycarboxylic acids. . . . .	362
4. Summary. . . . .	364
References. . . . .	365

\* Dedicated to the 60th birthday of Professor Dr. med. M. Eggstein.

## 1. INTRODUCTION

Diabetes mellitus is the most common of the metabolic diseases, its frequency in the general population being estimated to be more than 2%. It has no distinct and definable pathogenesis, etiology, invariable set of clinical findings and generally accepted laboratory tests. The disease is characterized by a number of hormone-induced metabolic abnormalities and by long-term complications.

Because abnormalities of glucose levels have been easiest to measure and were discovered first, diabetes mellitus has been primarily defined as a defect in the glucose metabolism. However, the metabolisms of lipids and amino acids are also affected. The development of the techniques for metabolite analysis, gas chromatographic-mass spectrometric (GC-MS) methods being the most flexible and probably the most powerful, has contributed to our understanding of the pathophysiological alterations in the glucose, lipid and amino acid metabolisms. Such alterations occur during situations of decompensation of the disease and diabetic ketoacidosis, but certain abnormalities are also found when diabetic patients are under good therapeutic control.

Diabetic patients are susceptible to an acute metabolic complication, viz., diabetic ketoacidosis. This profound alteration of metabolism results from a deficiency of insulin action. During diabetic ketoacidosis ketone bodies and four classes of organic acids, viz., fatty acids, dicarboxylic acids, oxocarboxylic acids and hydroxycarboxylic acids, are elevated in blood and in urine. They reflect increased lipolysis,  $\omega$ -oxidation of fatty acids, ketogenesis and degradation of amino acids, which are the causes of ketoacidosis. The levels of the metabolites normalize when the ketoacidotic situation is overcome.

Another organic acid, 2-ethyl-3-oxohexanoic acid, which is also an oxocarboxylic acid but behaves differently to the ketone bodies and the other oxocarboxylic acids, is increased in therapeutically well controlled diabetic patients.

Because of the variety of metabolic changes in diabetes mellitus, profile analyses of the metabolites are particularly suited to biochemical and diagnostic studies of the disease. With profile analyses it is possible to distinguish clearly between a metabolic situation where the diabetes is sufficiently balanced and a metabolic situation where the diabetes is decompensated. Certainly such information can also be obtained by measuring blood and urinary glucose, pH and the conventional ketone bodies, and this is done routinely. Profile analyses, however, give a much more detailed picture of the metabolic pathways involved, such as glucose metabolism only, or glucose metabolism and lipolysis, or in addition ketogenesis, or in addition metabolism of amino acids, or in addition lactic acidosis. Moreover, profile analyses supply information on the extent of the pathophysiological deviation and on the success of the therapeutic approach in rebalancing the diabetic patient.

Apart from this more diagnostically oriented application, profile analyses are helpful in biochemical studies of the metabolic pathways, e.g., in investigating if ketogenesis produces only acetoacetic acid, 3-hydroxybutyric acid and acetone, or other ketones as well. Profile analyses may also help to answer the question of whether the metabolism of a diabetic patient whose blood glucose is well controlled is otherwise normal.

## 2. PROFILING PROCEDURES

### 2.1. Methods of choice

Because of the complexity of the metabolites in blood and in urine, GC is the method of choice for the analysis of these compounds. High-performance liquid chromatography (HPLC) is less generally applicable because of the volatility of many of the aliphatic ketones. The factors which determine the separation efficiency, and the restrictions arising from these factors, are discussed in detail elsewhere.

This review covers methods of metabolite analysis obtained with GC and GC-MS.

### 2.2 Ketones

#### 2.2.1. Isolation of ketones

Of the neutral non-acidic ketone bodies, acetone, the most common, arises from ketogenesis. In addition to acetone, weight ketones occur in blood and urine, viz., methyl ketones.

The isolation of these ketones from biological fluids by adsorption on a solid material, such as Tenax GC, and alcohols, are also isolated and analyzed by GC profile analysis. Several years ago, the use of a polymer of 2,6-diphenyl-*p*-phenylene diamine as a method is applicable to serum and urine. The retention is 95–100% for ketones with increasing polarity of the substance. The adsorption on the Tenax GC is more carbon atoms. Strongly polar ketones are incompletely adsorbed (50–70%).

Other isolation techniques, such as solvent extraction, quantitative determinations of ketones are less suitable than the gas-phase method. Solvent impurities cause interference.

#### 2.2.2. Gas chromatographic profile analysis

For GC analysis, the compounds are adsorbed on a solid material. Usually thermal desorption is inserted into the GC injector. The injector is cooled pre-column prior to sample injection. Quantitative desorption of the ketones is achieved at 300°C. Substances with higher boiling points are not desorbed.

The GC separation can be achieved on a 10 m  $\times$  0.5 mm I.D. stainless-steel

## 2. PROFILING PROCEDURES

### 2.1. *Methods of choice*

Because of the complexity of the ketone bodies and the organic acids in blood and in urine, GC is the method of choice for the separation of these compounds. High-performance liquid chromatographic (HPLC) procedures are less generally applicable because of their limitations with regard to the detectability of many of the aliphatic components. Other disadvantages of HPLC are the separation efficiency, which is usually not adequate for organic acids, and the restrictions arising from coupling with MS.

This review covers methodical aspects and alternatives, and the results obtained with GC and GC-MS procedures, emphasizing profile analyses.

### 2.2 *Ketones*

#### 2.2.1. *Isolation of ketones*

Of the neutral non-acidic ketones, acetone is one of the three classical ketone bodies. It is formed by decarboxylation of acetoacetic acid, which itself arises from ketogenesis. In addition to acetone, a number of higher molecular weight ketones occur in blood and in urine. Like acetone, most of them are methyl ketones.

The isolation of these ketones can be achieved by gas-phase extraction and adsorption on a solid material. Other volatile endogenous constituents, e.g., alcohols, are also isolated and can be studied together with the ketones by profile analysis. Several years ago a number of variations of this isolation technique were suggested [1-7], most of them using Tenax GC, a porous polymer of 2,6-diphenyl-*p*-phenylene oxide, as the adsorbing material. The method is applicable to serum and urine. The efficiency of the gas-phase extraction is 95-100% for ketones with five or more carbon atoms. It decreases with increasing polarity of the substances and is only 50% for acetone. The extent of adsorption on the Tenax GC adsorbent is 95-100% for ketones with four or more carbon atoms. Strongly polar and hydrophilic substances such as acetone are incompletely adsorbed (50-80%).

Other isolation techniques, such as solvent extraction, have been applied in quantitative determinations of specific ketones. For profile analyses they are less suitable than the gas-phase extraction and adsorption procedure, because solvent impurities cause interferences in this low-molecular-weight range.

#### 2.2.2. *Gas chromatographic profiling of ketones*

For GC analysis, the components must be desorbed from the adsorbing material. Usually thermal desorption is chosen, the adsorbent trap being inserted into the GC injector block. The components are recondensed in a cooled pre-column prior to separation on a capillary GC column [4]. The quantitative desorption of the ketones requires a temperature of approximately 300°C. Substances with higher polarity are desorbed at 220°C and below.

The GC separation can be achieved on a mildly polar column, e.g., a 100 m × 0.5 mm I.D. stainless-steel column coated with Emulphor ON-870. The

column temperature was 60°C for 16 min, then programmed to 175°C at 2°C/min. Other stationary phases such as OV-17 and SE-30 can be used, as the separation of the ketones is not critical. The identification of the substances is performed by GC-MS.

Using the described profile analysis of volatile constituents in serum and urine, the ketones have been recognized as regular metabolites. In conjunction with the observation that the levels of the methyl ketones increase in diabetic or fasting ketoacidosis, these ketones have been identified as products arising from a general ketogenesis pathway. Another achievement of profile analysis of the volatile substances is the finding that aliphatic alcohols often reach elevated levels in the serum and urine of diabetic patients [8].

More insight into the pathophysiological situation has been obtained by complementing the profile analyses through quantitative determinations of the low-molecular-weight substances. For quantifications of ketones and alcohols, mass fragmentographic techniques have been developed [9-11].

### 2.3. Organic acids

#### 2.3.1. Isolation of organic acids

Two basic methods are used for the isolation of the organic acids: solvent extraction and isolation by anion-exchange chromatography. In solvent extraction, acceptable results have been obtained using diethyl ether [12, 13] or ethyl acetate [14-20], or a mixture of these solvents [21, 22]. Other solvents, such as dichloromethane [23] and hexane-diethyl ether (1:1) [24], have been less frequently used. To improve the extraction, the urine or serum sample can be saturated with salts, such as sodium chloride. The methods are not time consuming, as normally they are carried out batchwise and only in a few instances continuously. However, for many organic acids the procedure is not quantitative. This is especially true for polyhydroxycarboxylic acids and other very polar acids. Protein-containing fluids, especially serum, have to be deproteinated prior to extraction. The removal of the proteins can be achieved by dialysis or ultrafiltration [19, 25] or by precipitation [26]. As protein-bound organic acids are partially lost during dialysis or ultrafiltration, precipitation of the proteins is often preferable [27]. However, some organic acids may coprecipitate with the proteins.

Isolation of organic acids by anion-exchange chromatography is widely used for studying acid profiles. Usually DEAE-Sephadex [28-31], Amberlyst A-26 [32, 33] or Dowex [25] is employed. Isolation by anion exchange is more laborious than solvent extraction procedures. However, it is more quantitative, and it is chosen when detailed profiles of various classes of organic acids, including hydroxycarboxylic acids, are being investigated. The improvement in the isolation of polyhydroxycarboxylic acids achieved by using DEAE-Sephadex instead of diethyl ether-ethyl acetate has been described [34]. Good results for profiles have also been obtained on isolation using Extrelut columns [35].

#### 2.3.2. Derivatization of carboxyl groups

Generally, the carboxyl groups of organic acids are converted into either the

trimethylsilyl achieved with 31, 36-38] 39-41]. For reaction with advantages and mass spectra of methylsilyl est and more prec methylation w it does have t also methylate and methoxy possible reacti amino groups of the substanc control analys methane in the

Apart from tives of the car suggested.

In addition t can be derivati are converted methylsilylated methylated.

#### 2.3.3. Derivatization

The carbonyl many oxocarbo derivatives are O-ethyl oximes trimethylsilylqu

#### 2.3.4. Pre-fractionation

Because of th and in urine, fr is favoured. It b trations are to preparative thin HPLC [46].

In the investi mellitus, pre-fra using Kieselgel phenylacetate, n stances were app The plates were [n-heptane-diet.

trimethylsilyl esters or the methyl esters. Trimethylsilylation is usually achieved with bis(trimethylsilyl)trifluoroacetamide (BSTFA) [16, 18, 19, 21, 31, 36-38] or bis(trimethylsilyl)acetamide (BSA) [14, 15, 17, 20, 28, 39-41]. For the methylation of organic acids, most laboratories use the reaction with diazomethane [22, 25, 32, 33, 35, 42-47]. Both derivatives have advantages and disadvantages. When using GC-MS to identify the acids, the mass spectra of the methyl esters are easier to interpret than those of the trimethylsilyl esters, as the fragmentation of the methyl esters is often simpler and more predictable. When the necessary working precautions are followed, methylation with diazomethane is a very simple and rapid procedure. However, it does have the disadvantage that the hydroxy groups of phenolic acids are also methylated, making it impossible to distinguish between aromatic hydroxy and methoxy groups in the original compound. Another disadvantage is the possible reaction of diazomethane with double bonds, carbonyl groups and amino groups [48] to produce artifacts. To establish the original structure of the substances and to differentiate between genuine compounds and artifacts, control analyses with GC-MS have been performed using deuterated diazomethane in the sample preparation [47, 48].

Apart from trimethylsilyl and methyl esters, in some instances other derivatives of the carboxyl groups, such as pentafluorobenzyl esters [23], have been suggested.

In addition to the carboxyl group, other functional groups in organic acids can be derivatized. Whereas the hydroxy groups of hydroxycarboxylic acids are converted into the trimethylsilyl ethers when the organic acids are trimethylsilylated, they are usually left underivatized when the organic acids are methylated.

### 2.3.3. Derivatization of carbonyl groups

The carbonyl groups of oxocarboxylic acids are usually derivatized because many oxocarboxylic acids are not stable enough for GC analysis. Suitable derivatives are oximes [14, 17, 41, 49], O-methyloximes [18, 33, 50, 51] and O-ethyloximes [28, 37]. 2-Oxocarboxylic acids have been converted into O-trimethylsilylquinoxalins [52-54].

### 2.3.4. Pre-fractionation of organic acids

Because of the complexity of the mixture of acidic components in serum and in urine, frequently pre-fractionation of the total mixture of organic acids is favoured. It becomes essential whenever constituents present in low concentrations are to be studied. Pre-fractionation is effectively achieved either by preparative thin-layer chromatography (TLC) [32, 33, 46, 47, 50, 51] or by HPLC [46].

In the investigation of organic acids in samples from patients with diabetes mellitus, pre-fractionation of the acid derivatives by TLC into four fractions using Kieselgel 60 F<sub>254</sub> plates was applied [33]. Methyl stearate, methyl phenylacetate, methyl indolylbutyrate and dimethyl tartrate as reference substances were applied on the right-hand side of the plate to mark the fractions. The plates were developed with two solvent mixtures in succession. The first [*n*-heptane-diethyl ether-chloroform (40:30:30)] was allowed to run to the

upper edge of the plate. After marking the position of methyl indolylbutyrate under a UV lamp (254 nm), the plate was developed with the second solvent mixture [cyclohexane—chloroform—methanol (40:55:5)] up to the marked position of methyl indolylbutyrate. With the left-hand side of the plate covered, the right-hand side was treated with 2,7-dichlorofluorescein in order to make the reference substances visible at 366 nm.

Fraction 1 (zone between methyl stearate and methyl phenylacetate) contained mainly fatty acids, fraction 2 (zone between methyl phenylacetate and methyl indolylbutyrate) contained dicarboxylic acids, the derivatized oxocarboxylic acids and aromatic acids, fraction 3 (zone between methyl indolylbutyrate and dimethyl tartrate) contained hydroxycarboxylic acids, aromatic acids, nitrogen-containing acids and acid conjugates and fraction 4 (zone between dimethyl tartrate and the starting line) contained additional acid conjugates. For the analysis of the oxocarboxylic acids and the hydroxycarboxylic acids, further fractionation of fraction 2 in two subfractions (2a, 2b) and fraction 3 in four subfractions (3a–3d) gave more detailed information on these two classes of organic acids in diabetes mellitus [50].

An example of the analytical procedure for the organic acids, including pre-fractionation by preparative TLC, is shown in Fig. 1.




Fig. 1. Analytical procedure for the organic acids.

### 2.3.5. Gas chromatographic profiling of organic acids

The isolated and derivatized organic acids can be analysed by GC or GC-MS, either as a total profile or as partial profiles after pre-fractionation of the sample. Packed columns that were formerly used for the GC separation were usually replaced by glass or fused-silica capillary columns. Frequently used stationary phases include OV-17, OV-101, SE-30, OV-1701, DB-1 and DB-1701. The columns are usually 25 m long.

Total profiles of organic acids have been widely studied in conjunction with

the detection of metabolic pathways. In a biochemical study, an overall acid defect, caused by a metabolic defect, often leads to changes in the organic acid profile. Gas chromatography-mass spectrometry profiles have been used to identify organic acids [22, 33], such as phaeohydroxy acids in diabetic ketoacidosis. Capillary columns DB-1701 have been used for the analysis of programmed

The acid profiles. In an applied, such as acids and acidographic mode, the analysis of the O-methylated acids in their mass spectra in blood and urine (10 µg/dl and 10 µg/dl).

The advantage of this method, such as very small amount

It must be noted that artifact formation may be produced during the separation process. Many examples [40, 48, 55, 56]

## 3. APPLICATIONS

### 3.1. Diabetic profile

Profile analysis of therapeutically used organic acids, the ketone 4-heptanone. Because of the components, the 4-heptanone. The decarboxylation

the detection and characterization of pathophysiological situations and altered metabolic pathways leading to organic acidaemias and acidurias. Many of these biochemical alterations are sufficiently pronounced to be recognizable in the overall acid profile. Typical examples are inborn errors of metabolism. The defect, caused by an insufficient enzyme activity at a certain step of the metabolism, often of the metabolism of amino acids, results in specific and clear changes in the acid profile [55]. It has been proposed that screening for such organic acidurias can be achieved with GC alone and without mass spectrometry when a dual-column GC method is employed [17, 20]. Total profiles have also been used in the investigation of normal ranges of organic acids [22, 31], of abnormal levels of organic acids in patients with tumours such as phaeochromocytoma or neuroblastoma [56] and in the study of the metabolic changes in patients with diabetes mellitus, especially during diabetic ketoacidosis [57]. In the last example, 25 m × 0.2 mm I.D. glass capillary columns coated with OV-17 and fused-silica columns coated with OV-1701 have been used. The column temperature was 40°C for 10 min, then programmed to 230°C at 2°C/min.

The acid fractions are separated on the same stationary phases as the total profiles. In addition to flame ionization detectors, specific detectors have been applied, such as the thermionic specific detector for nitrogen-containing acids and acid derivatives, and the mass spectrometer in the mass fragmentographic mode. The thermionic specific detector has proved useful in the analysis of oxocarboxylic acids when the carbonyl function is converted into the O-methyloxime [33]. The identification of the organic acids is based on their mass spectra and on their GC retention indices. The levels of the different blood and urinary acids analysed by GC and identified by GC-MS are between 10 µg/dl and 100 mg/dl in serum or urine.

The advantages of the pre-fractionation procedure over the total profile method, such as improvement in separation and enrichment of acids present in very small amounts, have been described and exemplified [50].


It must be pointed out that in the analysis of organic acids the possibility of artifact formation as a source of error must be critically considered. Artifacts may be produced at any step of the sample preparation procedure and during the separation. Another source of error is interferences from drugs and medications. Many examples of artifacts and other interferences have been reported [40, 48, 55, 58-60].

### 3. APPLICATION TO DIABETES MELLITUS

#### 3.1. Diabetic patients under good therapeutic control

Profile analyses of serum and urine samples of diabetic patients who are therapeutically well controlled have shown that the levels of the ketones and the organic acids are not different from those of normal individuals, except for the ketone 4-heptanone and the organic acid 2-ethyl-3-oxohexanoic acid. Because of the chemical and biochemical relationship between these two components, the sum of 4-heptanone and 2-ethyl-3-oxohexanoic acid is called total 4-heptanone. The ketone is formed from the labile 3-oxocarboxylic acid by decarboxylation according to the reaction

C or GC-MS,  
ation of the  
paration were  
quently used  
DB-1 and  
unction with



The decarboxylation partly occurs *in vivo*, analogously to the decarboxylation of acetoacetic acid to form acetone. *In vitro*, heat enhances this reaction [9].

The acid is detected by GC-MS within a fraction of the organic acids either in the form of its methyl ester with the carboxyl group left underivatized, or in the form of its methyl ester with the carbonyl group derivatized to the O-methyl oxime [61, 62]. The ketone is detected within the profile of low-molecular-weight components by GC [2, 4, 62]. Total 4-heptanone is determined after thermal decarboxylation of the 3-oxocarboxylic acid [62].

The biochemical pathway of the formation of 2-ethyl-3-oxohexanoic acid is not completely clear. However, the findings suggest that the biosynthesis of the acid runs parallel to the synthesis of fatty acids, with butyryl-coenzyme A as the starter. This hypothesis is supported by the observation that together with elevated 2-ethyl-3-oxohexanoic acid, ethylmalonic acid, whose coenzyme A form is an intermediate of this mechanism, is also increased [63].

A study with 28 healthy individuals, 143 patients without metabolic diseases, 109 diabetic patients of type I and 125 diabetic patients of type II revealed increased levels of total 4-heptanone in the urine of the diabetic patients. Whereas the mean values for total urinary 4-heptanone in healthy individuals and non-diabetic patients were 207  $\mu\text{g}$  per 24 h and 246  $\mu\text{g}$  per 24 h, respectively, the mean urinary excretion of 2-ethyl-3-oxohexanoic acid plus 4-heptanone in the 234 diabetic patients was 1073  $\mu\text{g}$  per 24 h (diabetics of type I 1215  $\mu\text{g}$  per 24 h and diabetics of type II 987  $\mu\text{g}$  per 24 h) [63]. In a longitudinal study over 30 days, the total 4-heptanone values of two healthy controls were 80–410  $\mu\text{g}$  per 24 h.

It has also been observed that the urinary total 4-heptanone decreases during diabetic ketoacidosis, which is in accordance with the postulated biochemical origin of 2-ethyl-3-oxohexanoic acid. From the findings it is concluded that the excretion of total 4-heptanone is elevated in diabetes mellitus when the test is performed outside hyperglycaemic and ketoacidotic periods. It is further concluded that the excretion of total 4-heptanone is a sensitive indicator of the interplay between fatty acid oxidation and fatty acid synthesis. A decrease is observed during fatty acid oxidation and an increase during fatty acid synthesis [64].

### 3.2. Diabetic patients with ketoacidosis

#### 3.2.1. Ketones

In the profile of volatile metabolites a number of higher molecular weight methyl ketones are regularly found in addition to acetone. The main components are 2-pentanone, 3-penten-2-one and 2-heptanone (Fig. 2). During ketoacidosis their urinary excretion and their concentrations in serum are




Fig. 2. (a) Gas chromatogram of the metabolites in the urine of a patient with ketoacidosis. (b) Gas chromatogram of the metabolites in the urine of a patient with ketoacidosis. Column, 100 m  $\times$  0.5 mm; column temperature, 60°C for 16 min.

drastically increased. Acetone, the precursor, acetoacetic acid, i.e., 3-oxohexanoic acid, has [50].






Fig. 2. (a) Gas chromatogram of the methyl ketones in the urine of a normal individual. (b) Gas chromatogram of the methyl ketones in the urine of a patient with diabetic ketoacidosis. Column, 100 m  $\times$  0.5 mm I.D., stainless steel, coated with Emulphor ON-870; column temperature, 60°C for 16 min, then programmed to 175°C at 2°C/min.


drastically increased. Acetone is formed by spontaneous decarboxylation from its precursor, acetoacetic acid. The corresponding precursor of 2-pentanone, i.e., 3-oxohexanoic acid, has been found within a fraction of the organic acids [50].




Acetone                  Acetoacetic acid



2-Pentanone                  3-Oxohexanoic acid



3-Penten-2-one



2-Heptanone

Even though the precursors of 3-penten-2-one and 2-heptanone have not yet been detected, probably because of insufficient concentrations, a general concept for ketogenesis is likely. By analogy with the formation of the three conventional ketone bodies acetoacetic acid, 3-hydroxybutyric acid and




Fig. 5. Hypothetical general pathway for ketogenesis.

acetone, ov  
genesis path  
molecular w  
Under the  
ketone peak  
A comparis  
during ketoa

### 3.2.2. Fatty

The decre  
accumulatio  
3-hydroxybu  
into the uri  
of the incre  
the concentr  
serum. This  
class of acids

### 3.2.3. Alipha

Dicarboxy  
oxidation fo  
in a norm  
ketoacidosis  
20% of the f

A number  
are listed in  
The dicarbox

TABLE 1

### DICARBOXYL

No.	Acid
1	Malonic
2	Methyl
3	Ethylm
4	Succini
5	Methyls
6	2,3-Met
7	Glutaric
8	Ethylsu
9	3-Methy
10	3-Methy
11	2,3-Met
12	3-Methy
13	2-Ethyl
14	Adipic a
15	2-Methy
16	3-Methy
17	2,4-Dim

\* 3-Methylgluta

acetone, over the 3-hydroxy-3-methylglutaryl-CoA pathway, a general ketogenesis pathway over 3-hydroxy-3-alkylglutaryl-CoA would lead to the higher molecular weight ketone bodies (Fig. 3).

Under the experimental conditions for the profiles of volatile metabolites, a ketone peak represents the total ketone, which means ketone plus its precursor. A comparison of Fig. 2a and b also reveals a decrease in total 4-heptanone during ketoacidosis.

### 3.2.2. Fatty acids

The decreased pH of the blood during diabetic ketoacidosis is caused by an accumulation of organic acids, mainly fatty acids, acetoacetic acid and 3-hydroxybutyric acid. Whereas a large portion of the ketone bodies is excreted into the urine, fatty acids occur in urine only in small amounts. As a result of the increased lipolysis caused by the low insulin activity during ketoacidosis, the concentrations of the whole spectrum of free fatty acids are elevated in serum. This is a very familiar situation and therefore no emphasis is put on this class of acids.

### 3.2.3. Aliphatic dicarboxylic acids

Dicarboxylic acids are formed from long-chain monocarboxylic acids by  $\omega$ -oxidation followed by  $\beta$ -oxidation [65], which is a process of low priority in a normal metabolic situation, but which gains importance during ketoacidosis. It has been found in experiments with rats that between 5 and 20% of the fatty acids are metabolized by initial  $\omega$ -oxidation [66, 67].

A number of dicarboxylic acids have been identified in normal urine. Some are listed in Table 1 according to increasing methylene units on OV-1701. The dicarboxylic acids appear in fractions 2a and 2b of the TLC pre-fractiona-

TABLE 1

DICARBOXYLIC ACIDS IN NORMAL URINE

No.	Acid	No.	Acid
1	Malonic acid	18	Muconic acid
2	Methylmalonic acid	19	3,4-Methyleneadipic acid
3	Ethylmalonic acid	20	Pimelic acid
4	Succinic acid	21	3-Methylpimelic acid
5	Methylsuccinic acid	22	2,4-Dimethylpimelic acid
6	2,3-Methylenesuccinic acid	23	2,3-Methylenepimelic acid
7	Glutaric acid	24	3,4-Methylenepimelic acid
8	Ethylsuccinic acid	25	Suberic acid
9	3-Methylglutaric acid	26	3-Methylsuberic acid
10	3-Methylglutaconic acid*	27	2,3-Methylenesuberic acid
11	2,3-Methyleneglutaric acid	28	3,4-Methylenesuberic acid
12	3-Methylglutaconic acid*	29	Azelaic acid
13	2-Ethylglutaric acid	30	3-Methylazelaic acid
14	Adipic acid	31	3,4-Methyleneazelaic acid
15	2-Methyladipic acid	32	Sebacic acid
16	3-Methyladipic acid	33	5-Decylnedioic acid
17	2,4-Dimethyladipic acid	34	3,4-Methylenesebacic acid

\* 3-Methylglutaconic acid occurs in the form of the *cis-trans* isomers.




Fig. 4. Fraction 2b of the methyl esters of the organic acids in the urine of a normal individual. Column, 25 m  $\times$  0.2 mm I.D., fused silica, coated with OV-1701; column temperature 40°C for 10 min, then programmed to 230°C at 2°C/min. 3-Methylglutaconic acid occurs in the form of the *cis-trans* isomers.

tion. Fig. 4 shows fraction 2b of the organic acids in the urine of a normal individual, with some dicarboxylic acids labelled. Most of the other substances in fraction 2b are aromatic acids. In serum dicarboxylic acids are also found, but in much smaller concentrations than in urine. Therefore, some of the acids have not been detected so far. On the other hand, the higher molecular weight saturated and unbranched dicarboxylic acids ( $C_{12}$ – $C_{18}$ ) are identified in serum only.

During diabetic ketoacidosis mainly the saturated acids with even-numbered carbon chains are elevated, especially succinic, adipic, suberic and sebacic acids.




Fig. 5. Fraction 2b of the methyl esters of the organic acids in the urine of a diabetic patient with ketoacidosis. GC conditions as in Fig. 4.

The increase in succinic acid (Fig. 4). Suberic acid and sebacic acid are also elevated (Fig. 2a).

The methyl-branched dicarboxylic acids (with methyl groups on the ring) are little affected by ketoacidosis. Methylglutaric acid, methylglutaconic acid, 2,3-methylene-glutaric acid, 3-methylglutaric acid, 3-methylglutaconic acid, 3-methyladipic acid, 3,4-methyleneadipic acid, and 3-methylglutaconic acid are assumed to be products of the ketogenesis. This is supported by experiments with deuterated acetate. The methylglutaric acids are indeed of endogenous origin, as shown by the formation of diazomethane.

### 3.2.4. Aliphatic oxocarboxylic acids

Oxomonocarboxylic acids (with one or two methylene units) originate from the ketogenesis. They are produced from glucose in the liver. 3-Oxobutyric acid, caproic and 2-oxo-3-methylvaleric acid are products of the ketogenesis. 3-Oxobutyric acid is a product of ketogenesis according to the 3-hydroxy-3-methylglutaric acid, i.e., 3-oxobutyric acid, i.e., 3-oxobutyric acid, amounts in urine only during ketoacidosis, analogous to the ketogenesis of acetyl-CoA and acetyl-CoA as starting material. 3-Oxobutyric acid, which was found by Niemann in the urine of acidotic patients and was explained better understood as resulting from the ketogenesis of ethylhexanoic acid is assumed to be a product of the ketogenesis of 2-oxobutyric acid and 4-oxobutyric acid. This applies to 2-oxovaleric and 5-oxovaleric acid in the urine of ketoacidotic patients.

In the serum of normal individuals 3-oxobutyric acid, pyruvic and 2-oxoisocaproic acid are found in considerable but varying amounts. 3-Oxobutyric acid appears in the early portions of the urine.

In patients with ketoacidosis

TABLE 2

### OXOMONOCARBOXYLIC ACIDS

No.	Acid
1	Glyoxylic acid
2	Pyruvic acid
3	2-Oxobutyric acid
4	3-Oxobutyric acid
5	2-Oxoisovaleric acid
6	2-Oxovaleric acid

The increase in succinic acid and adipic acid is exemplified in Fig. 5 (cf., Fig. 4). Suberic acid and sebacic acid are not observed as components of fraction 2a.

The methyl-branched dicarboxylic acids and the acids with a cyclopropane ring are little affected by ketoacidosis. They are formed by  $\omega$ -oxidation of methyl-branched or methylene monocarboxylic acids [68, 69], which are assumed to be products of the microflora of the intestinal tract. It was shown by experiments with deuterated diazomethane that the methylene dicarboxylic acids are indeed of endogenous origin and are not artifacts produced with diazomethane.

#### 3.2.4. Aliphatic oxocarboxylic acids

Oxomonocarboxylic acids (listed in Table 2 according to increasing methylene units) originate from different biochemical sources. Pyruvic acid is produced from glucose in the glycolytic process, 2-oxoisovaleric, 2-oxoisocaproic and 2-oxo-3-methylvaleric acids stem from the oxidative transamination of the amino acids valine, leucine and isoleucine (Figs. 6-8). 3-Oxobutyric acid is a product of ketogenesis, which means that it is formed from acetyl-CoA according to the 3-hydroxy-3-methylglutaryl-CoA cycle. For the homologous 3-oxocarboxylic acid, i.e., 3-oxohexanoic acid, which occurs in detectable amounts in urine only during ketoacidosis, a condensation mechanism analogous to the ketogenesis pathway can be assumed (Fig. 3, with butyryl-CoA and acetyl-CoA as starters). The reduction product, 3-hydroxyhexanoic acid, which was found by Niwa et al. [19] in the urine and serum of ketoacidotic patients and was explained as an intermediate of  $\beta$ -oxidation, could be better understood as resulting from this condensation mechanism. 3-Oxo-2-ethylhexanoic acid is assumed to be formed according to a mechanism analogous to the fatty acid biosynthesis [63]. The formation of glyoxylic, 2-oxobutyric and 4-oxobutyric acids cannot be easily explained. The same applies to 2-oxovaleric and 5-oxohexanoic acids, which are often found in the urine of ketoacidotic patients.

In the serum of normal individuals the main oxomonocarboxylic acids are pyruvic and 2-oxoisocaproic acids [33]. In urine only pyruvic acid is excreted in considerable but varying amounts, the excretion of the other oxocarboxylic acids being low (Fig. 9). The derivatives of all of the oxomonocarboxylic acids appear in the early portions of the gas chromatograms of fractions 2a and 2b.

In patients with ketoacidosis, the ketone body 3-oxobutyric acid and all

TABLE 2  
OXOMONOCARBOXYLIC ACIDS IN NORMAL URINE

No.	Acid	No.	Acid
1	Glyoxylic acid	7	4-Oxobutyric acid
2	Pyruvic acid	8	2-Oxo-3-methylvaleric acid
3	2-Oxobutyric acid	9	2-Oxoisocaproic acid
4	3-Oxobutyric acid	10	3-Oxohexanoic acid
5	2-Oxoisovaleric acid	11	5-Oxohexanoic acid
6	2-Oxovaleric acid	12	3-Oxo-2-ethylhexanoic acid




Fig. 6. Oxo- and hydroxycarboxylic acids derived from valine.




Fig. 7. Oxo- and hydroxycarboxylic acids derived from leucine.




Fig. 8. Oxo- and hydroxycarboxylic acids derived from isoleucine.




Fig. 9. Early peaks in the urine chromatogram of a patient with maple syrup urine disease occur as symmetrical peaks.




Fig. 8. Oxo- and hydroxycarboxylic acids derived from isoleucine.




Fig. 9. Early portion of the gas chromatogram of fraction 2a of the derivatives of the organic acids in the urine of a normal individual. The O-methylxime derivatives of 2-oxoisovaleric acid occur as *syn-anti* isomers. GC conditions as in Fig. 4.




Fig. 10. Early portion of the gas chromatogram of fraction 2a of the derivatives of the organic acids in the urine of a diabetic patient with ketoacidosis. The O-methyloxime derivatives of 2-oxoisovaleric, 2-oxoisocaproic, 2-oxo-3-methylvaleric, 3-oxobutyric and 3-oxohexanoic (3-oxocaproic) acids occur as *syn-anti* isomers. GC conditions as in Fig. 4. (Reproduced from ref. 50.)

three 2-oxocarboxylic acids as metabolites of valine, leucine and isoleucine are raised [33]. The elevated level of the amino acid metabolites is not surprising as the branched-chain amino acids are also increased during ketoacidosis [70]. In urine a large amount of 3-oxobutyric acid is excreted. 2-Oxoisovaleric, 2-oxoisocaproic and 2-oxo-3-methylvaleric acids are also raised (Fig. 10). However, their absolute amounts in urine are low compared with serum. Apparently the 2-oxocarboxylic acids are reutilized and, in contrast to the ketone bodies, the organism loses only small amounts of these constituents.

### 3.2.5. Aliphatic hydroxycarboxylic acids

The hydroxymonocarboxylic acids detected in urine are listed in Table 3 according to increasing methylene units. With the exception of glycolic, 2-hydroxyisobutyric, 2-hydroxybutyric and 3-hydroxypropionic acids, the formation of which is not completely clarified, the biochemical origin of the other hydroxymonocarboxylic acids is easily traced. Lactic acid is produced by reduction of pyruvic acid under the conditions of anaerobic glycolysis. 3-Hydroxybutyric acid is formed by ketogenesis. The other hydroxymonocarboxylic acids are degradation products of valine, leucine and isoleucine. 2-Hydroxyisovaleric and 3-hydroxyisobutyric acids stem from valine (Fig. 6), 2-hydroxyisocaproic and 3-hydroxyisovaleric acids from leucine (Fig.

TABLE 3  
HYDROXYMONOCARBOXYLIC ACIDS

No.	Acid
1	Glycolic acid
2	Lactic acid
3	2-Hydroxyisobutyric acid
4	2-Hydroxybutyric acid
5	3-Hydroxypropionic acid
6	3-Hydroxyisovaleric acid
7	3-Hydroxybutyric acid

7) and 2-hydroxy-3-methylvaleric and 2-ethylpropionic acids from isoleucine is found as a diastereomeric mixture.

The hydroxymonocarboxylic acids are found [50]. Without the pre-fractionation cause some of them overlap in the chromatogram of ketoacidotic patients, where the 3-hydroxybutyric acid is the most prominent.

In patients with ketoacidosis the hydroxymonocarboxylic acid metabolites of the branched-chain amino acids show as examples the fractionation




Fig. 11. Gas chromatogram of fraction 3c from a normal individual. GC conditions as in Fig. 4.

TABLE 3  
HYDROXYMONOCARBOXYLIC ACIDS IN NORMAL URINE

No.	Acid	No.	Acid
1	Glycolic acid	8	2-Hydroxyisovaleric acid
2	Lactic acid	9	3-Hydroxyisobutyric acid
3	2-Hydroxyisobutyric acid	10	3-Hydroxy-2-methylbutyric acid
4	2-Hydroxybutyric acid	11	2-Hydroxy-3-methylvaleric acid
5	3-Hydroxypropionic acid	12	2-Hydroxyisocaproic acid
6	3-Hydroxyisovaleric acid	13	3-Hydroxy-2-ethylpropionic acid
7	3-Hydroxybutyric acid		

7) and 2-hydroxy-3-methylvaleric, 3-hydroxy-2-methylbutyric and 3-hydroxy-2-ethylpropionic acids from isoleucine (Fig. 8). 3-Hydroxy-2-methylbutyric acid is found as a diastereomeric pair, the *threo*-form being the main diastereomer.

The hydroxymonocarboxylic acids appear in the four subfractions 3a–3d [50]. Without the pre-fractionation their identification would be difficult, because some of them overlap in the GC separation, especially with samples from ketoacidotic patients, where 3-hydroxybutyric acid is by far the major component.

In patients with ketoacidosis, 3-hydroxybutyric acid and all the hydroxy metabolites of the branched-chain amino acids are elevated. Figs. 11 and 12 show as examples the fractions 3c of the hydroxymonocarboxylic acids in the




Fig. 11. Gas chromatogram of the methyl esters of the hydroxycarboxylic acids in fraction 3c from a normal individual. GC conditions as in Fig. 4. (Reproduced from ref. 50.)




Fig. 12. Gas chromatogram of the methyl esters of the hydroxycarboxylic acids in fraction 3c from a diabetic patient with ketoacidosis. GC conditions as in Fig. 4. (Reproduced from ref. 50.)

urine of a normal person and a ketoacidotic patient, respectively. The remainder of the hydroxycarboxylic acids are found in fractions 3a, 3b and 3d. It is observed that the 3-hydroxy series is particularly affected by ketoacidosis. Of the two possible metabolites of the 3-hydroxy series resulting from isoleucine, 3-hydroxy-2-ethylpropionic acid is the major product. For 3-hydroxy-2-methylbutyric acid an elevation is not always found. The three components of the 2-hydroxy series of the amino acid metabolites are increased. However, their absolute amounts remain low. In normal urines 2-hydroxyisocaproic acid is often not even detectable. The behaviour of the hydroxymonocarboxylic acids in serum appears to be similar to that in urine.

#### 4. SUMMARY

Diabetes mellitus is a defect not only in glucose metabolism, but also in the metabolism of lipids and amino acids. Gas chromatographic and gas chromatographic-mass spectrometric profile analyses have contributed much to the understanding of the metabolic changes connected with this defect. Ketones are isolated by a gas-phase extraction and adsorption technique and profiled after thermal desorption. Organic acids are isolated by solvent extraction or anion exchange, derivatized and separated either as total acid profiles or sub-profiles after pre-fractionation of the acid derivatives.

The main results are as follows: (a) In patients with diabetes mellitus, especially in those therapeutically well controlled, the concentration of ketone bodies is elevated to higher molecular weight ketone bodies. (c) During diabetes mellitus the following acids are elevated: 3-hydroxybutyric acid, resulting from  $\omega$ - and  $\beta$ -oxidation of long-chain carboxylic acids as metabolites of amino acids and as products of ketogenesis originating from amino acids and

#### REFERENCES

- 1 A. Zlatkis, W. Bertsch, H.A. Bartsch, G. Coscia and N. Fleischer, *Anal. Biochem.* 1967, 21, 100.
- 2 M. Novotny, M.L. McConnell, *J. Clin. Invest.* 1967, 42, 100.
- 3 A. Zlatkis, W. Bertsch, D.A. Bartsch, *J. Clin. Invest.* 1967, 42, 100.
- 4 H.M. Liebich and O. Al-Babbili, *J. Clin. Invest.* 1967, 42, 100.
- 5 A. Zlatkis and K. Kim, *J. Clin. Invest.* 1967, 42, 100.
- 6 H.M. Liebich and J. Wöll, *J. Clin. Invest.* 1967, 42, 100.
- 7 A. Zlatkis, K.Y. Lee, C.F. Poo, *J. Clin. Invest.* 1967, 42, 100.
- 8 H.M. Liebich, O. Al-Babbili, *J. Clin. Invest.* 1967, 42, 100.
- 9 H.M. Liebich and G. Huesgen, *J. Clin. Invest.* 1967, 42, 100.
- 10 H.M. Liebich, *J. Chromatogr.* 1967, 1, 100.
- 11 H.M. Liebich, H.J. Bülow and E. Jellum, *J. Clin. Invest.* 1967, 42, 100.
- 12 E. Jellum, O. Stokke and L. Eide, *J. Clin. Invest.* 1967, 42, 100.
- 13 J.E. Pettersen and E. Jellum, *J. Clin. Invest.* 1967, 42, 100.
- 14 H.J. Sternowsky, J. Roboz, F. P. O'Connell, *J. Clin. Invest.* 1967, 42, 100.
- 15 T. Niwa, T. Ohki, K. Maeda, *J. Clin. Invest.* 1967, 42, 100.
- 16 J. Greter, S. Lindstedt, H. Seer, *J. Clin. Invest.* 1967, 42, 100.
- 17 K. Tanaka, A. West-Dull, *J. Clin. Invest.* 1967, 42, 100.
- 18 M.L. Holland, G.R. Rhodes, D. A. Zlatkis, *J. Clin. Invest.* 1967, 42, 100.
- 19 T. Niwa, K. Yamada, T. Ohki and M. Y. Tsai, *J. Clin. Invest.* 1967, 42, 100.
- 20 M.Y. Tsai, C. Oliphant and M. L. Holland, *J. Clin. Invest.* 1967, 42, 100.
- 21 W.L. Fitch, P.J. Anderson and H.M. Liebich, *J. Clin. Invest.* 1967, 42, 100.
- 22 H.M. Liebich, A. Pickert, U. Stöckel, *J. Clin. Invest.* 1967, 42, 100.
- 23 I. Penttilä, A. Huhtikangas, *J. Clin. Invest.* 1967, 42, 100.
- 24 L.R. Alexander and J.B. Justice, *J. Clin. Invest.* 1967, 42, 100.
- 25 H. Thoma, J. Reiner and G. Spittler, *J. Clin. Invest.* 1967, 42, 100.
- 26 H.M. Liebich, A. Pickert and E. Jellum, *J. Clin. Invest.* 1967, 42, 100.
- 27 H.M. Liebich, A. Pickert, C. Oliphant, *J. Clin. Invest.* (1984) 118.
- 28 R.A. Chalmers and R.W.E. Waller, *J. Clin. Invest.* 1967, 42, 100.
- 29 R.A. Chalmers, A.M. Lawson and R.W.E. Waller, *J. Clin. Invest.* 1967, 42, 100.
- 30 A.M. Lawson, R.A. Chalmers and R.W.E. Waller, *J. Clin. Invest.* 1967, 42, 100.
- 31 S.C. Gates, N. Dendramis and M. Spittler, *J. Clin. Invest.* 1967, 42, 100.
- 32 M. Spittler and G. Spittler, *J. Clin. Invest.* 1967, 42, 100.
- 33 H.M. Liebich, A. Pickert and J. Wöll, *J. Clin. Invest.* 1967, 42, 100.
- 34 P. Sims, R. Truscott and B. Hall, *J. Clin. Invest.* 1967, 42, 100.
- 35 D. Pinkston, G. Spittler, H. Thoma, *J. Clin. Invest.* 1967, 42, 100.

The main results are as follows. (a) Increased total 4-heptanone is inherently connected with diabetes mellitus. Its urinary levels are elevated in therapeutically well controlled patients. (b) A general ketogenesis pathway leads to higher molecular weight ketone bodies in addition to the conventional ketone bodies. (c) During diabetic ketoacidosis, in addition to the fatty acids the following acids are elevated in serum and in urine: dicarboxylic acids resulting from  $\omega$ - and  $\beta$ -oxidation of monocarboxylic acids; oxomonocarboxylic acids as metabolites of the amino acids valine, leucine and isoleucine and as products of ketogenesis; and hydroxymonocarboxylic acids, also originating from amino acids and from ketogenesis.

## REFERENCES

- 1 A. Zlatkis, W. Bertsch, H.A. Lichtenstein, A. Tishbee, F. Shunbo, H.M. Liebich, A.M. Coscia and N. Fleischer, *Anal. Chem.*, 45 (1973) 763.
- 2 M. Novotny, M.L. McConnell, M.L. Lee and R. Farlow, *Clin. Chem.*, 20 (1974) 1105.
- 3 A. Zlatkis, W. Bertsch, D.A. Bafus and H.M. Liebich, *J. Chromatogr.*, 91 (1974) 379.
- 4 H.M. Liebich and O. Al-Babbili, *J. Chromatogr.*, 112 (1975) 539.
- 5 A. Zlatkis and K. Kim, *J. Chromatogr.*, 126 (1976) 475.
- 6 H.M. Liebich and J. Wöll, *J. Chromatogr.*, 142 (1977) 505.
- 7 A. Zlatkis, K.Y. Lee, C.F. Poole and G. Holzer, *J. Chromatogr.*, 163 (1979) 125.
- 8 H.M. Liebich, O. Al-Babbili, A. Zlatkis and K. Kim, *Clin. Chem.*, 21 (1975) 1294.
- 9 H.M. Liebich and G. Huesgen, *J. Chromatogr.*, 126 (1976) 465.
- 10 H.M. Liebich, *J. Chromatogr.*, 112 (1975) 551.
- 11 H.M. Liebich, H.J. Bülow and R. Kallmayr, *J. Chromatogr.*, 239 (1982) 343.
- 12 E. Jellum, O. Stokke and L. Eldjarn, *Clin. Chem.*, 18 (1972) 800.
- 13 J.E. Pettersen and E. Jellum, *Clin. Chim. Acta*, 41 (1972) 199.
- 14 H.J. Sternowsky, J. Roboz, F. Hutterer and G. Gaull, *Clin. Chim. Acta*, 47 (1973) 371.
- 15 T. Niwa, T. Ohki, K. Maeda, A. Saito and K. Kobayashi, *Clin. Chim. Acta*, 99 (1979) 71.
- 16 J. Greter, S. Lindstedt, H. Seemann and G. Steen, *Clin. Chem.*, 26 (1980) 261.
- 17 K. Tanaka, A. West-Dull, D.G. Hine, T.B. Lynn and T. Loewe, *Clin. Chem.*, 26 (1980) 1847.
- 18 M.L. Holland, G.R. Rhodes, D. Wiesler and M. Novotny, *J. Chromatogr.*, 306 (1984) 23.
- 19 T. Niwa, K. Yamada, T. Ohki and H. Furukawa, *J. Chromatogr.*, 337 (1985) 1.
- 20 M.Y. Tsai, C. Oliphant and M.W. Josephson, *J. Chromatogr.*, 341 (1985) 1.
- 21 W.L. Fitch, P.J. Anderson and D.H. Smith, *J. Chromatogr.*, 162 (1979) 249.
- 22 H.M. Liebich, A. Pickert, U. Stierle and J. Wöll, *J. Chromatogr.*, 199 (1980) 181.
- 23 I. Penttilä, A. Huhtikangas, J. Herranen and O. Moilanen, *J. Chromatogr.*, 338 (1985) 265.
- 24 L.R. Alexander and J.B. Justice, Jr., *J. Chromatogr.*, 342 (1985) 1.
- 25 H. Thoma, J. Reiner and G. Spiteller, *J. Chromatogr.*, 309 (1984) 17.
- 26 H.M. Liebich, A. Pickert and B. Tetschner, *J. Chromatogr.*, 289 (1984) 259.
- 27 H.M. Liebich, A. Pickert, C. Först, B. Tetschner and T. Risler, *Krankenhausarzt*, 57 (1984) 118.
- 28 R.A. Chalmers and R.W.E. Watts, *Analyst (London)*, 97 (1972) 958.
- 29 R.A. Chalmers, A.M. Lawson and R.W.E. Watts, *Clin. Chim. Acta*, 52 (1974) 43.
- 30 A.M. Lawson, R.A. Chalmers and R.W.E. Watts, *Clin. Chem.*, 22 (1976) 1283.
- 31 S.C. Gates, N. Dendramis and C.C. Sweeley, *Clin. Chem.*, 24 (1978) 1674.
- 32 M. Spiteller and G. Spiteller, *J. Chromatogr.*, 164 (1979) 253.
- 33 H.M. Liebich, A. Pickert and J. Wöll, *J. Chromatogr.*, 217 (1981) 255.
- 34 P. Sims, R. Truscott and B. Halpern, *J. Chromatogr.*, 222 (1981) 337.
- 35 D. Pinkston, G. Spiteller, H. von Henning and D. Matthaëi, *J. Chromatogr.*, 223 (1981) 1.

- 36 C. Jakobs, E. Solem, J. Ek, K. Halvorsen and E. Jellum, *J. Chromatogr.*, 143 (1977) 31.  
 37 R.A. Chalmers, A.M. Lawson and O. Borud, *Clin. Chim. Acta*, 77 (1977) 117.  
 38 T. Niwa, K. Maeda, T. Ohki and J. Sakakibara, *J. Chromatogr.*, 228 (1982) 59.  
 39 F.W. Bultitude and S.L. Newham, *Clin. Chem.*, 21 (1975) 1329.  
 40 M. Duran, D. Ketting, S.K. Wadman, C. Jakobs, R.B.H. Schutgens and H.A. Veder, *Clin. Chim. Acta*, 90 (1978) 187.  
 41 K. Tanaka, D.G. Hine, A. West-Dull and T.B. Lynn, *Clin. Chem.*, 26 (1980) 1839.  
 42 J.E. Pettersen, E. Jellum and L. Eldjarn, *Clin. Chim. Acta*, 38 (1972) 17.  
 43 L. Borg, S. Lindstedt and G. Steen, *Clin. Chim. Acta*, 41 (1972) 363.  
 44 S. Landaas, *Clin. Chim. Acta*, 54 (1974) 39.  
 45 S. Lindstedt, K. Norberg, G. Steen and E. Wahl, *Clin. Chem.*, 22 (1976) 1350.  
 46 A. Grupe and G. Spittler, *J. Chromatogr.*, 226 (1981) 301.  
 47 H.M. Liebich and A. Pickert, *J. Chromatogr.*, 338 (1985) 25.  
 48 H.M. Liebich and C. Först, *J. Chromatogr.*, 338 (1985) 33.  
 49 R.A. Chalmers and R.W.E. Watts, *Analyst (London)*, 97 (1972) 951.  
 50 H.M. Liebich and C. Först, *J. Chromatogr.*, 309 (1984) 225.  
 51 H.M. Liebich and C. Först, *J. Chromatogr.*, 338 (1985) 187.  
 52 U. Langenbeck, A. Hoinowski, K. Mantel and H.-U. Möhring, *J. Chromatogr.*, 143 (1977) 39.  
 53 U. Langenbeck, A. Mench-Hoinowski, K.-U. Dieckmann, H.-U. Möhring and M. Petersen, *J. Chromatogr.*, 145 (1978) 195.  
 54 R.J. Early, J.R. Thompson, T. McAllister, T.W. Fenton and R.J. Christopherson, *J. Chromatogr.*, 310 (1984) 1.  
 55 E. Jellum, *J. Chromatogr.*, 143 (1977) 427.  
 56 S.C. Gates, C.C. Sweeley, W. Krivit, D. de Witt and B.E. Blaisdell, *Clin. Chem.*, 24 (1978) 1680.  
 57 H.M. Liebich, in Lu Pei-chang and E. Bayer (Editors), *Proceedings of Sino-West German Symposium on Chromatography*, Science Press, Beijing, China, 1983, pp. 195ff.  
 58 C. Jakobs, M. Bojasch, M. Duran, D. Ketting, S.K. Wadman and D. Leupold, *Clin. Chim. Acta*, 106 (1980) 85.  
 59 M. Ende, P. Pfeifer and G. Spittler, *J. Chromatogr.*, 183 (1980) 1.  
 60 S. Bauer, M. Neupert and G. Spittler, *J. Chromatogr.*, 309 (1984) 243.  
 61 H.M. Liebich, G. Huesgen, A. Pickert, U. Stierle and J. Wöll, *J. High Resolut. Chromatogr. Chromatogr. Commun.*, 6 (1983) 507.  
 62 H.M. Liebich, *J. Chromatogr.*, 273 (1983) 67.  
 63 H.M. Liebich, U. Stierle and M. Eggstein, *Verh. Dtsch. Ges. Inn. Med.*, 89 (1983) 698.  
 64 H.M. Liebich and M. Eggstein, poster presented at the XII International Congress of Clinical Chemistry, Rio de Janeiro, 1984.  
 65 J.E. Pettersen, *Clin. Chim. Acta*, 41 (1972) 231.  
 66 I. Björkhem, *J. Lipid Res.*, 19 (1978) 585.  
 67 W. Kam, K. Kumaran and B.R. Landau, *J. Lipid Res.*, 19 (1978) 519.  
 68 J.E. Pettersen and O. Stokke, *Biochim. Biophys. Acta* 304 (1973) 316.  
 69 J. Greter, S. Lindstedt and G. Steen, *J. Biol. Chem.*, 254 (1979) 2807.  
 70 S.A. Adibi, *Metabolism*, 25 (1976) 1287.

*Journal of Chromatography  
Biomedical Applications*  
Elsevier Science Publishers

CHROMBIO. 2989

REVIEW

MICROBIAL CHEMISTRY

CHROMATOGRAPHY  
TECHNIQUES

ILIA BRONDZ\*

*Department of Chemistry*  
and

INGAR OLSEN

*Department of Microbiology*

(First received October 1984)

CONTENTS

1. Introduction. . . . .
  - 1.1. Microbial taxonomy. . . . .
  - 1.2. Chemotaxonomy. . . . .
2. Bacterial products and their analysis. . . . .
  - 2.1. Volatile and metabolites. . . . .
  - 2.2. Extracts. . . . .
  - 2.3. Whole defatted cells. . . . .
  - 2.4. Whole cells. . . . .
  - 2.5. Single colony analysis. . . . .
    - 2.5.1. Pyrolysis-mass spectrometry. . . . .
3. Cell envelope components. . . . .
  - 3.1. Sugars. . . . .
  - 3.2. Teichoic/teichuroic acids. . . . .
  - 3.3. Lipids and their analysis. . . . .
    - 3.3.1. Sample preparation. . . . .
    - 3.3.2. Gas chromatography. . . . .
    - 3.3.3. Mass spectrometry. . . . .
    - 3.3.4. High-performance liquid chromatography. . . . .
    - 3.3.5. Thin-layer chromatography. . . . .
  - 3.4. Lipopolysaccharides. . . . .
  - 3.5. Murein. . . . .

0378-4347/86/\$03.50